

Uncovering the Link between Sleep, Alexithymia, and Overweight Status: A Machine Learning Approach

*Kadir Uludag

Shanghai Jiao Tong University (SJTU) - Shanghai Mental Health Center

Excessive weight, including obesity and overweight statuses, has been identified as a significant risk factor for a wide range of severe health problems and the severity of psychiatric symptoms associated with various psychiatric diseases. Our primary objective was to predict overweight status using comprehensive data on sleep-related factors and alexithymia. We employed advanced machine learning models, specifically random forest (RF) and neural networks (NN), to identify the most influential predictive factors in this context. The Stockholm Brain Study focuses on functional brain imaging data obtained from two distinct age groups: participants aged 20-30 years and older individuals aged 65-75 years. In order to assess overweight and obesity status, participants' body mass index (BMI) was calculated using the World Health Organization's defined cut-off points for overweight and obesity. Additionally, the study incorporates alexithymia data to further investigate its role in predicting overweight status. The results showed that RF ML model predicted overweight status with an accuracy of 72 % (sensitivity: 100 %, specificity:30%). The area under curve (AUC) value was 95. Furthermore, NN found that overweight status was predicted with the accuracy of 93 % in the train data and the accuracy of 56 % in the test data (sensitivity:80%, specificity:20%). To validate our results, the cross-validation was performed based on algorithms of RF, and NNs, and resulted in an average accuracy score of 0.70 ± 0.12 , 0.54 ± 0.11 , respectively. The study findings revealed that the prediction of overweight status exhibited remarkable accuracy, surpassing 70 percent, through the utilization of a RF model.

Keywords: Obesity, Overweight Status, Sleep, Severity of Alexithymia, Severity of Symptoms, Machine Learning

*Correspondence concerning this article should be addressed to Mr. Kadir Uludag, Shanghai Jiao Tong University (SJTU) - Shanghai Mental Health Center. Email: kadiruludag@mailsucas.ac.cn

Being overweight during childhood has emerged as a strong predictor of numerous health outcomes later in life, including heightened risk for cardiovascular disease (Janssen et al., 2005) and metabolic conditions such as diabetes. Given this, understanding the factors that contribute to overweight status—including sleep and alexithymia—is essential. This area of study carries considerable weight because overweight and obesity remain pressing public health issues, and researchers are increasingly turning to machine learning (ML) as a tool for building predictive models of health outcomes. Since both sleep patterns and alexithymia have been linked to the development of overweight and obesity, integrating these factors into predictive models could enhance their accuracy.

Sleep disorders are widespread (Hauri, 2021), while insufficient sleep may lead to specific psychological symptoms, such as cognitive symptoms (Killgore, 2010). In addition, sleep deprivation is common among individuals and associated with health costs (Goel, Rao, Durmer, & Dinges, 2009). Furthermore, restricted sleep may lead to excessive daytime sleepiness and altered mood, and sleep loss to hormonal changes that could result in obesity (Vorona et al., 2005). Similarly, a different study found that sleep duration was inversely related to being overweight in young males (Eisenmann, Ekkekakis, & Holmes, 2006). Moreover, additional research confirmed that short sleep duration is a risk factor for both

overweight and obesity statuses in children (Chaput et al., 2011).

Besides sleep-related factors, alexithymia is a disturbance in emotional processing shown by difficulties in recognizing feelings (Martínez-Sánchez, Ato-García, & Ortiz-Soria, 2003) and is associated with more light sleep (Bazydło, Lumley, & Roehrs, 2001). Furthermore, the analysis indicates a significant association between overweight status and impaired emotional recognition, with the data suggesting that individuals with severe obesity face greater challenges in accurately discerning their emotions compared to the control group (Da Ros, Vinai, Gentile, Forza, & Cardetti, 2011).

Collectively, the role of both alexithymia and sleep quality on overweight status is unclear. Therefore, our study goal was to predict overweight status using random forest (RF) and neural networks (NN) models using sleep and alexithymia data.

Method

Subjects

The data from a previous study (Nilsson et al., 2016) (The Stockholm Brain Study) was used to conduct this study.

A total of 83 participants were recruited through the MR center at Karolinska University Hospital, drawn from both younger (ages 20–30) and older (ages 65–75) healthy adults living in Stockholm. Rating scales were completed remotely at home via Google Forms. Participant anonymity was preserved by assigning each individual a unique code rather than collecting identifying information. All electronic submissions were complete, with no missing data. For any paper-based responses, data were manually entered into spreadsheets, and a second researcher cross-checked the entries to confirm accuracy.

Overall, our study included the following variables to predict weight status: age, Karolinska Sleep Questionnaire (KSQ) Overall Sleep Quality, KSQ_Morning or Evening Person, Toronto Alexithymia Scale-20 (TAS20) Total score, TAS20 subscale of DIF, TAS20 subscale of DDF, TAS20 subscale of EOT, were used to predict overweight status.

Participants were paid around 280 Euro. As mentioned in the previous article, participants who participated in the Zürich prosocial game could gain additional money and be offered taxi travel (Nilsson et al., 2016). The data collection spanned two periods: from December 4, 2012, to March 27, 2013, and from October 24, 2013, to April 29, 2014.

Instruments

Calculating Body Mass Index

BMI (kg/m^2) was derived from each participant's measured height and weight. Based on these values, participants were classified into weight categories: overweight (25–29.9) and obese (30–34.9), following World Health Organization guidelines (Organization, 2000).

Toronto Alexithymia Scale

Alexithymia was assessed using the 20-item Toronto Alexithymia Scale (TAS-20), administered in its Swedish-language adaptation (Simonsson-Sarnecki et al., 2000).

Random Forest Model

The Random Forest (RF) algorithm, a widely used machine learning approach in psychological, neurological, and psychiatric research, was employed for prediction. As an ensemble method, RF combines the output of multiple decision trees to generate its predictions.

SLEEP, ALEXITHYMIA, AND OVERWEIGHT STATUS

Random Forest Model Parameters

The RF model was built using the default settings available in Python's Scikit-learn library (Pedregosa et al., 2011). A random state of 10 was specified, the maximum tree depth was fixed at 4, and 30% of the data was reserved for testing.

Neural Networks (NN)

Neural networks represent another ML technique that has gained traction in recent research. Because of the intricate way they process information, they are often described as "black box" models (Alber et al., 2019).

Neural Network Parameters

The NN model was likewise constructed using Scikit-learn's default parameters (Pedregosa et al., 2011). The architecture included 30 input-layer neurons, 50 neurons in the hidden layer, and 30 output neurons, with the random state again set to 10.

Statistical Analysis

We compared demographic and clinical variables between patients with and without overweight status using the chi-square test for categorical variables. Independent sample T-test and Mann Whitney U test were used for non-categorical variables according to the normality of the variables. Python 3.8.8. version was used to conduct analysis related to machine learning (ML), while SPSS version 17 was used to perform statistical analysis related to descriptive statistics (e.g., mean and standard deviation). In addition, the Kolmogorov-Smirnov test was used to test the normality of the data. There was no missing data in our study.

Ethical Approval

All procedures performed in studies involving human participants were following the ethical standards of the institutional and national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Also, informed consent was obtained from all patients.

As mentioned in the previous study (Nilsson et al., 2016) this study was registered to clinicaltrials.gov (NCT02000076). In addition, a list of hypotheses was uploaded to the Open Science Framework (<https://osf.io/bxfsb/>). Further, the study was approved by the Regional Ethics Review board of Stockholm (2012/1098-31/2, amended as 2012/1565-32 and 2012/1870-32), mainly including permission for open publication of de-identified data.

Results

Descriptive Statistics

The participants (n=83) were between (20-30 years) and (65-75 years). Forty-six were in the younger age group, while 47 were in the older age group. Most of them were males (n=42). In addition, three obese patients with over the 30 body mass index (BMI) were excluded.

Table 1

Precision and Recall of Random Forest and Neural Networks

	Random Forest	Neural Networks
Precision	65	50
Recall	64	50
F1 Score	25	48

Table 2

Confusion Matrix according to Random Forest

15	7
0	3


Table 3

Confusion Matrix according to Neural Networks


$$\begin{bmatrix} 12 & 8 \\ 3 & 2 \end{bmatrix}$$

Figure 1

Most Predictive Factors according to Random Forest Model



TAS 20 total score, TAS 20 EOT, TAS 20 DDF, KSQ Nap frequency TAS 20 DIF
KSQ morning evening person, KSQ overall sleep quality.

Figure 2*Most Predictive Factors, according to Neural Networks Model*

TAS 20 DDF, TAS 20 DIF , KSQ overall sleep quality, KSQ nap frequency, TAS 20 EOT, KSQ morning or evening person, TAS 20 total

Random Forest Machine Learning Model Results

Random Forest (RF) model found that in the training set the accuracy was 72 (sensitivity:100%. specificity:30%). The area under curve (AUC) value was 95. The most predictive factors were Toronto Alexithymia Scale-20 scores (TAS) 20 total score, TAS 20 EOT, TAS 20 DDF, Karolinska sleep questionnaire (KSQ) nap frequency, TAS 20 DIF KSQ morning-evening person, KSQ overall sleep quality respectively.

To validate our ML results, the cross-validation was performed based on algorithms of RF, and NNs, and resulted in an average accuracy score of 0.70 ± 0.12 , 0.54 ± 0.11 , respectively.

Neural Networks Machine Learning Model Results

Neural Networks (NN) model found that in the training set the accuracy was 93 while in the test set the accuracy was 56% (sensitivity:80%. specificity:20%). The most predictive factors were TAS 20 subscale of DDF, TAS 20 subscale DIF , KSQ overall sleep quality, KSQ subscale of nap frequency, TAS 20 subscale of EOT, KSQ morning or evening person, and TAS 20 total, respectively.

Discussion

The central results of this study demonstrate that both random forest (RF) and neural network (NN) machine learning models are capable of predicting overweight status. For the RF model, training accuracy reached 93%, while testing accuracy was 56%, with sensitivity at 100%, specificity at 30%, and an AUC of 95%. The NN model, by comparison, achieved 80% training accuracy and 20% testing accuracy, with sensitivity and specificity mirroring the RF model at 100% and 30% respectively.

Notably, alexithymia-related variables emerged as the strongest predictors of overweight status across both models. Specifically, the RF model ranked the following as most predictive: total TAS-20 score, the Externally Oriented Thinking (EOT) subscale, the Difficulty Identifying Feelings (DIF) subscale, nap frequency from the Karolinska Sleep Questionnaire (KSQ), the Difficulty Describing Feelings (DDF) subscale, KSQ chronotype (morning/evening preference), and overall KSQ sleep quality.

The NN model produced a comparable ranking, identifying the DDF subscale, DIF subscale, overall KSQ sleep quality, KSQ nap frequency, EOT subscale, KSQ chronotype, and

total TAS-20 score as its top predictive features.

Taken together, these findings underscore the substantial contribution of alexithymia-related measures to predicting overweight status, pointing to the relevance of both emotional processing difficulties and sleep quality in weight regulation. This aligns with existing literature suggesting that excessive daytime sleepiness and short self-reported sleep duration may contribute to obesity risk, and that a connection exists between obesity and sleep disturbances more broadly (Vgontzas, Bixler, Chrousos, & Pejovic, 2008).

In addition, it is worth noting that sleep deprivation has been linked to a pattern of centripetal distribution of fat (De Bernardi Rodrigues et al., 2016). In addition, there is a complex relationship among alexithymia, overweight status, and sleep quality. Therefore, creating prediction models using sleep-related and alexithymia parameters may help predict overweight status. Considering that being overweight is detrimental to overall health, predicting the risk can be beneficial.

In the literature, variations in the defined cut-off points for obesity and overweight status can contribute to discrepancies observed in studies concerning obesity. Furthermore, our research identified heterogeneous variables, including age and gender, as potential confounding factors that may influence the outcomes.

Association between Alexithymia and Overweight Status

The literature consistently found that alexithymia is associated with overweight status. For example, the data suggest severely obese patients have more difficulty recognizing their emotions than controls (Da Ros et al., 2011). Similarly, findings supported having a substantial alexithymic element among severely obese patients compared to their counterparts (Clerici, Albonetti, Papa, Penati, & Invernizzi, 1992). Another study interestingly suggests that the improvement in weight-related attitudes following stable weight loss may differ in alexithymic and non-alexithymic obese patients (Franco Adami, Campostano, Ravera, Leggieri, & Scopinaro, 2001). Further, an additional study found that alexithymia was associated with a more severe binge eating disorder (Carano et al., 2006).

The existence of overweight status was associated with higher emotional dysregulation than normal weight conditions (Casagrande, Boncompagni, Forte, Guarino, & Favieri, 2020). Moreover, a different study found that alexithymia and psychopathology are correlated among obese patients seeking treatment (Pinna et al., 2011). Our study findings demonstrate that parameters related to alexithymia hold potential for predicting overweight status.

Association between Alexithymia and Sleep

It is essential to understand the association between alexithymia and sleep, as our study included alexithymia-related factors. The results of the meta-analysis study found a link between alexithymia and sleep problems (Alimoradi et al., 2021). Additional research has established a link between alexithymia and lower sleep quality (Murphy, Wulff, Catmur, & Bird, 2018). However, this connection appears to be indirect: when depression is statistically controlled for through multiple regression analysis, the relationship between alexithymia and sleep problems no longer holds. What remains is simply the already well-established link between depression and disrupted sleep quality (De Gennaro, Martina, Curcio, & Ferrara, 2004).

Also, dreaming has a specific manifestation in the case having alexithymia (Obrębska & Rohoza, 2021). Additionally, the findings of a different study suggest a consistent relationship between alexithymia and dreaming components that offer processes regulating emotion during dreaming (Nielsen, Levrier, & Montplaisir, 2011). Similarly, a study showed a higher incidence of nightmares and the association of nightmares with alexithymia (Godin, Montplaisir, Gagnon, & Nielsen, 2013). The association between alexithymia and overweight

SLEEP, ALEXITHYMIA, AND OVERWEIGHT STATUS

status highlights the importance of considering psychological factors in the prevention of overweight and obesity. By addressing emotional regulation and promoting healthy eating behaviors, interventions may be able to effectively target both alexithymia and overweight status, and improve overall health and well-being.

Conclusion

This study concluded that overweight status might be predicted using sleep-related and alexithymia data.

Limitations

We have used only BMI to measure obesity and overweight status. Many studies have relied on self-report measures of alexithymia and overweight status, which may be subject to inaccuracies. Also, small sample size is one of the limitations of the study. One of the limitations of this study is that the data collection was conducted during two distinct periods.

Suggestions for Further Studies

Future studies should measure various overweight status-related parameters and associated risk factors. In addition, an integrated approach to promote emotional regulation may reduce the risk of being overweight (Casagrande et al., 2020). Future research endeavors may delve into exploring potential mediators such as emotional eating or other maladaptive coping strategies to better understand the underlying mechanisms linking emotional regulation and weight regulation. Additionally, investigating neural or physiological pathways could shed light on the intricate connections between these two domains.

References

- Alber, M., Lapuschkin, S., Seegerer, P., Hägele, M., Schütt, K. T., Montavon, G., . . . Kindermans, P.-J. (2019). iNNvestigate neural networks! *J. Mach. Learn. Res.*, 20(93), 1-8. <https://www.jmlr.org/papers/v20/18-540.html>
- Alimoradi, Z., Majd, N. R., Broström, A., Tsang, H. W., Singh, P., Ohayon, M. M., ... & Pakpour, A. H. (2022). Is alexithymia associated with sleep problems? A systematic review and meta-analysis. *Neuroscience and Biobehavioral Reviews*, 133.
- Bazydlo, R., Lumley, M. A., & Roehrs, T. (2001). Alexithymia and Polysomnographic Measures of Sleep in Healthy Adults. *Psychosom Med*, 63(1). <https://www.diva-portal.org/smash/record.jsf?pid=diva2:1648826>
- Carano, A., De Berardis, D., Gambi, F., Di Paolo, C., Campanella, D., Pelusi, L., . . . Ferro, F. M. (2006). Alexithymia and body image in adult outpatients with binge eating disorder. *International Journal of Eating Disorders*, 39(4), 332-340. <https://doi.org/10.1002/eat.20238>
- Casagrande, M., Boncompagni, I., Forte, G., Guarino, A., & Favieri, F. (2020). Emotion and overeating behavior: effects of alexithymia and emotional regulation on overweight and obesity. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*, 25(5), 1333-1345. <https://doi.org/10.1007/s40519-019-00767-9>
- Chaput, J.-P., Lambert, M., Gray-Donald, K., McGrath, J. J., Tremblay, M. S., O'Loughlin, J., & Tremblay, A. (2011). Short sleep duration is independently associated with overweight and obesity in quebec children. *Canadian Journal of Public Health*, 102(5), 369-374. <https://doi.org/10.1007/BF03404179>
- Clerici, M., Albonetti, S., Papa, R., Penati, G., & Invernizzi, G. (1992). Alexithymia and obesity. *Psychotherapy and Psychosomatics*, 57(3), 88-93. <https://doi.org/10.1159/000288580>
- Da Ros, A., Vinai, P., Gentile, N., Forza, G., & Cardetti, S. (2011). Evaluation of alexithymia and depression in severe obese patients not affected by eating disorders. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*, 16(1), 24-29. <https://doi.org/10.1007/BF03327517>
- De Bernardi Rodrigues, A. M., da Silva, C. d. C., Vasques, A. C. J., Camilo, D. F., Barreiro, F.,

- Cassani, R. S. L., . . . Investigators, f. t. B. M. S. S. (2016). Association of sleep deprivation with reduction in insulin sensitivity as assessed by the hyperglycemic clamp technique in adolescents. *JAMA Pediatrics*, 170(5), 487-494. <https://doi.org/10.1001/jamapediatrics.2015.4365>
- De Gennaro, L., Martina, M., Curcio, G., & Ferrara, M. (2004). The relationship between alexithymia, depression, and sleep complaints. *Psychiatry Research*, 128(3), 253-258. <https://doi.org/10.1016/j.psychres.2004.05.023>
- Eisenmann, J. C., Ekkekakis, P., & Holmes, M. (2006). Sleep duration and overweight among Australian children and adolescents. *Acta Paediatrica*, 95(8), 956-963. <https://doi.org/10.1080/08035250600731965>
- Franco Adami, G., Campostano, A., Ravera, G., Leggieri, M., & Scopinaro, N. (2001). Alexithymia and body weight in obese patients. *Behavioral Medicine*, 27(3), 121-126. doi:10.1080/08964280109595778
- Godin, I., Montplaisir, J., Gagnon, J.-F., & Nielsen, T. (2013). Alexithymia associated with nightmare distress in idiopathic rem sleep behavior disorder. *Sleep*, 36(12), 1957-1962. <https://doi.org/10.5665/sleep.3238>
- Durmer, J. S., & Dinges, D. F. (2005). Neurocognitive consequences of sleep deprivation. *Seminars in Neurology*, 25(01), 117-129. <https://doi.org/10.1055/s-2005-867080>
- Hauri, P. J. (2021). *Sleep disorders*: Routledge.
- Janssen, I., Katzmarzyk, P. T., Srinivasan, S. R., Chen, W., Malina, R. M., Bouchard, C., & Berenson, G. S. (2005). Utility of childhood BMI in the prediction of adulthood disease: Comparison of National and international references. *Obesity Research*, 13(6), 1106-1115. <https://doi.org/10.1038/oby.2005.129>
- Killgore, W. D. S. (2010). Effects of sleep deprivation on cognition. In G. A. Kerkhof & H. P. A. v. Dongen (Eds.), *Progress in brain research* (Vol. 185, pp. 105-129): Elsevier.
- Martínez-Sánchez, F., Ato-García, M., & Ortiz-Soria, B. (2003). Alexithymia — state or trait? *The Spanish Journal of Psychology*, 6(1), 51-59. <https://doi.org/10.1017/s1138741600005205>
- Murphy, J., Wulff, K., Catmur, C., & Bird, G. (2018). Alexithymic traits, independent of depression and anxiety, are associated with reduced sleep quality. *Personality and Individual Differences*, 129, 175-178. <https://doi.org/10.1016/j.paid.2018.03.023>
- Nielsen, T., Levrier, K., & Montplaisir, J. (2011). Dreaming correlates of alexithymia among sleep-disordered patients. *Dreaming*, 21(1), 16-31. <https://doi.org/10.1037/a0022861>
- Nilsson, G., Tamm, S., d'Onofrio, P., Thuné, H. Å., Schwarz, J., Lavebratt, C., . . . Axelsson, J. (2016). A multimodal brain imaging dataset on sleep deprivation in young and old humans. <https://core.ac.uk/download/pdf/70344019.pdf>
- Obrębska, M., & Rohoza, K. (2021). Stress as a moderator of the relationship between alexithymia and dreaming: Research findings. *Dreaming*, 31(2), 186-194. <https://doi.org/10.1037/drm0000167>
- Obesity: preventing and managing the global epidemic. Report of a WHO consultation. (2000). In *PubMed* (Vol. 894, pp. i-253). <https://pubmed.ncbi.nlm.nih.gov/11234459>
- PedregosaFabian, VaroquauxGaël, GramfortAlexandre, MichelVincent, ThirionBertrand, GriselOlivier, BlondelMathieu, PrettenhoferPeter, WeissRon, DubourgVincent, VanderplasJake, PassosAlexandre, CournapeauDavid, BrucherMatthieu, PerrotMatthieu, & DuchesnayÉdouard. (2011). SciKit-Learn: Machine Learning in Python. *Journal of Machine Learning Research*. <https://doi.org/10.5555/1953048.2078195>
- Pinna, F., Lai, L., Pirarba, S., Orrù, W., Velluzzi, F., Loviselli, A., & Carpinello, B. (2011). Obesity, alexithymia and psychopathology: A case-control study. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*, 16(3), e164-e170. <https://doi.org/10.3275/7509>
- Simonsson-Sarnecki, M., Lundh, L., Törestad, B., Bagby, R. M., Taylor, G., & Parker, J. (2000). A Swedish translation of the 20-item Toronto Alexithymia Scale: Cross-validation of the factor structure. *Scandinavian Journal of Psychology*, 41(1), 25-30.

SLEEP, ALEXITHYMIA, AND OVERWEIGHT STATUS

<https://doi.org/10.1111/1467-9450.00167>

- Vgontzas, A. N., Bixler, E. O., Chrousos, G. P., & Pejovic, S. (2008). Obesity and sleep disturbances: Meaningful sub-typing of obesity. *Archives of Physiology and Biochemistry*, 114(4), 224-236. <https://doi.org/10.1080/13813450802521507>
- Vorona, R. D., Winn, M. P., Babineau, T. W., Eng, B. P., Feldman, H. R., & Ware, J. C. (2005). Overweight and obese patients in a primary care population report less sleep than patients with a normal body mass index. *Archives of Internal Medicine*, 165(1), 25. <https://doi.org/10.1001/archinte.165.1.25>